

# PRE-ANESTHESIA TESTING GRID

	Patient Type	ECG	CBC	BMP	POCT Glucose	PT/INR	HepACT	LFT	T&S	Urine HCG
PROCEDURE TYPE	I	NO PREOPERATIVE TESTING REQUIRED								
	II	TESTING BASED ON PATIENT COMORBIDITIES								
	III		X						X	
	IV	X	X	X		X		X	X	
CARDIAC COMORBIDITIES	CAD, Chest pain, CHF, Dysrhythmias, MI, Pacemaker/AICD, H/O Coronary stents/PTCA, Valvular disease, Previous cardiac surgery	X								
	Hyperlipidemia	X <sup>1</sup>								
	Hypertension	X <sup>1</sup>		X						
	Pulmonary hypertension	X								
	Vascular disease (peripheral or cerebral)	X		X						
OTHER COMORBID CONDITIONS	Anemia, bleeding disorder		X							
	BMI > 35				X <sup>2</sup>					
	Chronic kidney disease		X	X						
	Diabetes	X <sup>3</sup>		X	X <sup>2</sup>					
	End stage liver disease, Cirrhosis		X	X		X <sup>2</sup>	X <sup>2</sup>	X		
	End stage renal disease, Dialysis dependant	X	X	X <sup>2</sup>						
MEDICATIONS	ACEI, ARB, Digoxin, Diuretics			X						
	Anticoagulants		X <sup>4</sup>							
	Low-Molecular Wt. Heparin		X				X <sup>2</sup>			
	Intraop IV contrast			X						
	Warfarin		X			X <sup>2</sup>				
OTHER	All females 12-51									X <sup>2,5</sup>
	Age > 75		X	X						

LAB: Results good for 3 months (without clinical change)

ECG: Results good for 6 months (without clinical change)

<sup>1</sup> Age > 50

<sup>2</sup> Day of surgery

<sup>3</sup> Type I or on Insulin (Age >40); Type II (Age >50)

<sup>4</sup> All anticoagulants, except Aspirin

<sup>5</sup> Exceptions include previous hysterectomy, current pregnancy, miscarriage or ectopic